arXiv:1212.3031v2 [hep-th] 8 Jan 2013

KIAS-P12059

Nonextremal Kerr/CFT on a stretched horizon

Ee Chang-Young[1,*] and Myungseok Eune[2,†]

[1] *Department of Physics and Institute of Fundamental Physics,*
Sejong University, Seoul, 143-747, Korea
[2] *Research Institute for Basic Science,*
Sogang University, Seoul, 121-742, Korea

(Dated: January 9, 2013)

Abstract

We study the conformal symmetry described by $SL(2,R)_L \times SL(2,R)_R$ in the nonextremal Kerr black hole case. We calculate the central charges $c_{L,R}$ separately on a stretched horizon by the Hamiltonian gravity. In order to get two sets of conformal symmetry, we consider two Killing vectors each of which depends on one of the two independent coordinates separated from the angular coordinate of the corotating frame at the horizon. This was done in the spirit of point-splitting but with asymmetric separations. Then with appropriate choice of parameters, we obtain the Bekenstein-Hawking entropy by the Cardy formula.

---

[*] cylee@sejong.ac.kr

[†] younms@sogang.ac.kr

I. INTRODUCTION

Since the successful description of the holographic duality between the geometry near the horizon of extremal Kerr black hole and the conformal field theory [1–5], there have been many studies for the holographic duality in various black holes [6–23]. The holographic duality has been also developed for near extremal black holes by introducing the extremal parameter [15, 17] or calculating the graybody factor [24–27]. And the hidden conformal field theory described by the $SL(2,R)_L \times SL(2,R)_R$ in the Kerr black hole has been studied from the scalar wave equation [28]. However, in this case the values of the central charges $c_{L,R}$ have been just assumed to be the same as in the extremal black hole case. Up to now, the left and right central charges $c_{L,R}$ have been calculated only in the near extremal case [15]. There have been many extended studies of the hidden conformal symmetry for various black holes [29–41], but no calculation of the both central charges has been done. Recently, for both extremal and nonextremal Kerr black holes, the central charge has been obtained successfully by using the Hamiltonian gravity on a stretched horizon, but it describes only $SL(2,R)_L$ symmetry, not $SL(2,R)_L \times SL(2,R)_R$ [42–46]. There also have been studies for extremal black holes using the Hamiltonian gravity [47, 48].

We will investigate the conformal symmetry described by $SL(2,R)_L \times SL(2,R)_R$ on a stretched horizon in the nonextremal Kerr black hole case, adopting the method of Carlip [45, 46] and obtaining the central charges $c_{L,R}$ separately. In Sec. II, we review briefly the Hamiltonian gravity. In Sec. III, we obtain the central charges $c_{L,R}$, the temperatures $T_{L,R}$, and the total entropy S on a stretched horizon in the nonextremal case with the Hamiltonian gravity. In order to have two Virasoro algebras we consider two Killing vectors which are functions of either one of the two coordinates split from the angular coordinate of the corotating frame at the horizon. This splitting was done in the spirit of point-splitting, but with asymmetric separations from the angular coordinate. Then the central charges and the total entropy are evaluated. Finally, in Sec. IV, we conclude with discussion.

II. BRIEF REVIEW ON HAMILTONIAN GRAVITY

In order to obtain the central charges, we evaluate the central terms in the Hamiltonian gravity in this work. Here, we briefly review the related ingredients in the Hamiltonian gravity for this calculation. The detailed review can be found in Ref. [45]. We consider a stationary nonextremal black hole, whose metric can be written in the ADM form as follows.

$$ds^2 = -N^2 dt^2 + q_{ij}(dx^i + N^i dt)(dx^j + N^j dt). \tag{1}$$

Under a diffeomorphism generated by a vector field ξ^μ, the metric (1) transforms as

$$\delta_\xi N = \bar{\partial}_t \xi^\perp + \hat{\xi}^i \partial_i N, \tag{2}$$

$$\delta_\xi N^i = \bar{\partial}_t \hat{\xi}^i - N q^{ij} \partial_j \xi^\perp + q^{ik} \partial_k N \xi^\perp + \hat{\xi}^j \partial_j N^i, \tag{3}$$

$$\delta_\xi q_{ij} = q_{ik} \left(\partial_j \hat{\xi}^k - \frac{\partial_j N^k}{N} \xi^\perp \right) + q_{jk} \left(\partial_i \hat{\xi}^k - \frac{\partial_i N^k}{N} \xi^\perp \right) + \frac{1}{N} \xi^\perp \bar{\partial}_t q_{ij} + \hat{\xi}^k \partial_k qij, \tag{4}$$

where the convective derivative $\bar{\partial}_t$ is defined by

$$\bar{\partial}_t \equiv \partial_t - N^i \partial_i, \tag{5}$$

and the Killing vector ξ^μ is redefined as

$$\xi^\perp \equiv N \xi^t, \qquad \hat{\xi}^i \equiv \xi^i + N^i \xi^t, \tag{6}$$

which are called the "surface deformaton parameters". The symmetries of canonical general relativity are generated by the Hamiltonian

$$H[\xi] = \int d^3x \left(\xi^\perp \mathcal{H} + \hat{\xi}^i \mathcal{H}_i \right), \tag{7}$$

with

$$\mathcal{H} = \frac{1}{\sqrt{q}} (\pi^{ij} \pi_{ij} - \pi^2) - \sqrt{q} \mathcal{R}, \qquad \mathcal{H}^i = -2 D_j \pi^{ij}, \tag{8}$$

where q_{ij}, π^{ij}, and $\mathcal{R}$ are the spatial metric, the canonical momentum, and the spatial curvature scalar, respectively, and D_i is the spatial covariant derivative with respect to q_{ij}. $\mathcal{H}$ and $\mathcal{H}^i$ are the Hamiltonian and momentum constraints, respectively. In Refs. [45, 49], a new generator with a well-defined variation and with no boundary terms is defined by

$$\bar{H}[\xi] = H[\xi] + B[\xi], \tag{9}$$

where $B[\xi]$ depends only on the fields and parameters at the boundary, and is chosen to cancel the boundary terms in the variation of $H[\xi]$. The Poisson bracket of $\bar{H}[\xi]$ can be written as

$$\{\bar{H}[\xi], \bar{H}[\eta]\} = \bar{H}[\{\xi, \eta\}_{SD}] + K[\xi, \eta], \tag{10}$$

where the central term is given by

$$K[\xi,\eta] = B[\{\xi,\eta\}_{SD}] - \frac{1}{8\pi G}\int_{\partial\Sigma} d^2x\sqrt{\sigma}n^k\left[\frac{1}{\sqrt{q}}\pi_{ik}\{\xi,\eta\}^i_{SD} - \frac{1}{2\sqrt{q}}(\hat{\xi}_k\eta^\perp - \hat{\eta}_k\xi^\perp)\mathcal{H}\right.$$
$$+ (D_i\hat{\xi}_k D^i\eta^\perp - D_i\hat{\eta}_k D^i\xi^\perp) - (D_i\hat{\xi}^i D_k\eta^\perp - D_i\hat{\eta}^i D_k\xi^\perp)$$
$$\left. + \frac{1}{\sqrt{q}}\left(\hat{\eta}_k\pi^{mn}D_m\hat{\xi}_n - \hat{\xi}_k\pi^{mn}D_m\hat{\eta}_n\right)\right]. \tag{11}$$

Here, the surface deformaton brakets are given by

$$\{\xi,\eta\}^\perp_{SD} = \hat{\xi}^i D_i\eta^\perp - \hat{\eta}_i D_i\xi^\perp,$$
$$\{\xi,\eta\}^i_{SD} = \hat{\xi}^i D_k\hat{\eta}^i - \hat{\eta}^k D_k\hat{\xi}^i + q^{ik}\left(\xi^\perp D_k\eta^\perp - \eta^\perp D_k\xi^\perp\right). \tag{12}$$

We can calculate the central charges from the central term $K[\xi,\eta]$ given in Eq. (11). When a Virasoro subalgebra of the group of surface deformation is found in the following form, $\{\xi,\eta\}_{SD} = \xi\eta' - \eta\xi'$ where the prime denotes the derivative with respect to the rotating angle φ, then the boundary contribution $B[\{\xi,\eta\}_{SD}]$ can be ignored for our purpose [45]. This is because that the central charge of the Virasoro algebra is given by the coefficient of the following expression, $\int d\varphi(\xi'\eta'' - \eta'\xi'')$ that we may get from K in Eq. (11).

III. CFT IN NONEXTREMAL ROTATING BLACK HOLE

We now consider the nonextremal Kerr black hole governed by

$$ds^2 = -\frac{\Delta}{\Sigma}\left(dt - a\sin^2\theta\, d\varphi\right)^2 + \frac{\Sigma}{\Delta}dr^2 + \Sigma\, d\theta^2 + \frac{\sin^2\theta}{\Sigma}\left[adt - (r^2+a^2)d\varphi\right]^2, \tag{13}$$

in the Boyer-Lindquist coordinates, with

$$\Delta(r) = r^2 + a^2 - 2Mr, \tag{14}$$

$$\Sigma(r,\theta) = r^2 + a^2\cos^2\theta, \tag{15}$$

where M is the mass of the black hole and a is a parameter related to the angular momentum J of the black hole as $J = aM$. The nonextremal Kerr black hole (13) can be written in the ADM form,

$$ds^2 = -N^2dt^2 + d\rho^2 + q_{\varphi\varphi}(d\varphi + N^\varphi dt)^2 + q_{\theta\theta}d\theta^2, \tag{16}$$

where ρ is the proper distance from the horizon. In general, the line element of the nonextremal stationary rotating black hole can be expressed as in Eq. (16). Thus, from now on we consider the nonextremal stationary rotating black hole in four dimensions. Near the horizon of the black hole, the lapse function N and the shift vector N^φ in the metric (16) can be expanded as

$$N = \kappa_H \rho + \frac{1}{3!}\kappa_2(\theta)\rho^3 + \cdots , \tag{17}$$

$$N^\varphi = -\Omega_H - \frac{1}{2}\omega_2(\theta)\rho^2 + \cdots , \tag{18}$$

$$q_{\varphi\varphi} = q^{(H)}_{\varphi\varphi}(\theta) + \frac{1}{2}q^{(2)}_{\varphi\varphi}(\theta)\rho^2 + \cdots , \tag{19}$$

$$q_{\theta\theta} = q^{(H)}_{\theta\theta}(\theta) + \frac{1}{2}q^{(2)}_{\theta\theta}(\theta)\rho^2 + \cdots , \tag{20}$$

where κ_H and Ω_H are the surface gravity and the angular velocity on the event horizon, respectively. For convenience, we introduce the parameter ε defined by $\varepsilon \equiv -\frac{1}{2}\omega_2(\theta)\rho^2$ so that we have $N^\varphi \approx -\Omega_H + \varepsilon$ up to the order ρ^2. The shift vector N^φ becomes the angular velocity near horizon, when $\rho \to 0$.

When the black hole geometry is given by Eq. (16), the diffeomorphism given by (2)-(4) becomes

$$\xi^\rho = -\rho\bar{\partial}_t\xi^t, \tag{21}$$

$$\bar{\partial}_t\xi^\rho = \kappa_H^2\rho^2\partial_\rho\xi^t, \tag{22}$$

$$\bar{\partial}_t\hat{\xi}^\varphi = \kappa^2\rho^2 q^{\varphi\varphi}\partial_\varphi\xi^t - \frac{2\varepsilon}{\rho}\xi^\rho, \tag{23}$$

$$\partial_\rho\hat{\xi}^\varphi = -q^{-1}_{\varphi\varphi}\partial_\varphi\xi^\rho + \frac{2\varepsilon}{\rho}\xi^t, \tag{24}$$

where the convective derivative $\bar{\partial}_t$ in Eq. (5) becomes

$$\bar{\partial}_t = \partial_t - N^\varphi\partial_\varphi \approx \partial_t + (\Omega_H - \varepsilon)\partial_\varphi. \tag{25}$$

We choose the stretched horizon as a surface with a slightly different angular velocity $\bar{\Omega} \equiv \Omega_H - \bar{\varepsilon}$, where it is assumed that $\bar{\varepsilon}$ is proportional to ρ, where as the subleading term ε of the angular velocity near the horizon is proportional to ρ^2. For the purpose of studying the conformal field theory, we introduce a new coordinate system defined by

$$\tilde{t} = (\varphi - \bar{\Omega}t)/\Omega_H, \tag{26}$$

$$\tilde{\varphi} = \alpha[\varphi - (\Omega_H + \varepsilon)t], \tag{27}$$

dubbed as the right and the left chiral coordinates, respectively. Here, α is a constant that we shall determine later. The two coordinates $\tilde{t}$ and $\tilde{\varphi}$ are basically proportional to

$\varphi - (\Omega_H - \bar{\varepsilon})t$ and $\varphi - (\Omega_H + \varepsilon)t$, respectively, which are slightly separated from the angular coordinate of the corotating frame at the horizon, $\bar{\varphi} = \varphi - \Omega_H t$. Note that $\bar{\varepsilon}$ and ε are the order of ρ and ρ^2, respectively. We may consider these two coordinates as twofold splits of the angular coordinate of the corotating frame at the horizon in the spirit of point-splitting [50], but with asymmetric separations. Then, we assume the existence of two Killing vectors which are functions of either $x_R \equiv \varphi - \bar{\Omega}t$ or $x_L \equiv \varphi - (\Omega_H + \varepsilon)t$, and we dub them the right and left handed Killing vectors, respectively.

Now we first consider the right handed conformal field theory (RCFT) where the Killing vector ξ_R is a function of $x_R = \varphi - \bar{\Omega}t$. In this case, the expressions for the diffeomorphism invariance given by Eqs. (21), (22), (23), and (24) become

$$\xi_R^\rho = -\bar{\varepsilon}\rho\partial_\varphi\xi_R^t, \tag{28}$$

$$\bar{\varepsilon}\partial_\varphi\xi_R^\rho = \kappa^2\rho^2\partial_\rho\xi_R^t, \tag{29}$$

$$\bar{\varepsilon}\partial_\varphi\hat{\xi}_R^\varphi = q^{\varphi\varphi}\kappa_H^2\rho^2\partial_\varphi\xi_R^t - \frac{2\varepsilon}{\rho}\xi_R^\rho, \tag{30}$$

$$\partial_\rho\hat{\xi}_R^\varphi = -q_{\varphi\varphi}^{-1}\partial_\varphi\xi_R^\rho + \frac{2\varepsilon}{\rho}\xi_R^t, \tag{31}$$

up to the leading order ρ. From Eq. (11), the terms that contribute to the central charge is given by

$$K[\xi_R, \eta_R] = -\frac{\bar{\varepsilon}^3}{8\pi G q^{\varphi\varphi}\kappa_H^3\rho^2}\cdot\frac{A}{2\pi}\int d\varphi\left(\partial_\varphi\hat{\xi}_R^\varphi\partial_\varphi^2\hat{\eta}_R^\varphi - \partial_\varphi\hat{\eta}_R^\varphi\partial_\varphi^2\hat{\xi}_R^\varphi\right), \tag{32}$$

where A denotes the area of the stretched horizon, and the central charge can be read as

$$c_R = \frac{3\bar{\varepsilon}^3 A}{\pi G q^{\varphi\varphi}\kappa_H^3\rho^2}. \tag{33}$$

Next, we consider the left handed conformal field theory (LCFT) where the Killing vector ξ_L is a function of $x_L = \varphi - (\Omega_H + \varepsilon)t$. In this case, we have $\bar{\partial}_t\xi_L = -2\varepsilon\partial_\varphi\xi_L$. Then, the diffeomorphism invariance (3) can be written as

$$\delta_{\xi_L}N^\varphi \approx -2\varepsilon\partial_\varphi\hat{\xi}_L^\varphi + \frac{2\varepsilon}{\rho}\hat{\xi}_L^\rho = 0, \tag{34}$$

and we get

$$\hat{\xi}_L^\rho = \rho\partial_\varphi\hat{\xi}_L^\varphi. \tag{35}$$

Since the only nonvanishing component of the canonical momentum π^{ij} near the horizon is given by [45]

$$\pi^{\rho\varphi} = -\frac{\omega_2}{2\kappa_H}\sqrt{q} + \mathcal{O}(\rho^2), \tag{36}$$

the relevant terms for the central charge remained in K, given by Eq. (11), are

$$\begin{aligned} K[\xi,\eta] &= -\frac{1}{8\pi G}\int_{\partial\Sigma} d^2x \frac{\sqrt{\sigma}}{\sqrt{q}} n^k(\hat{\eta}_k \pi^{mn} D_m \hat{\xi}_n - \hat{\xi}_k \pi^{mn} D_m \hat{\eta}_n) \\ &= -\frac{\omega_2}{16\pi G\kappa_H}\int d^2x\sqrt{\sigma}\left(\hat{\xi}^\rho \partial_\varphi \hat{\eta}^\rho - \hat{\eta}^\rho \partial_\varphi \hat{\xi}^\rho\right). \end{aligned} \tag{37}$$

Plugging Eq. (35) into the above expression, we get

$$K[\xi_L, \eta_L] = \frac{\varepsilon}{8\pi G\kappa_H}\cdot\frac{A}{2\pi}\int d\varphi\left(\partial_\varphi \hat{\xi}^\varphi_L \partial^2_\varphi \hat{\eta}^\varphi_L - \partial_\varphi \hat{\eta}^\varphi_L \partial^2_\varphi \hat{\xi}^\varphi_L\right), \tag{38}$$

and the central charge can be read as

$$c_L = \frac{3\varepsilon A}{\pi G\kappa_H}. \tag{39}$$

For a scalar field in the Frolov-Thorne vacuum, the wave function in the eigenmodes of energy ω and angular momentum m can be written as $\Phi \sim e^{-i\omega t + im\varphi}$. Then, using the relation

$$e^{-i\omega t + im\varphi} = e^{-in_R \tilde{t} + in_L \tilde{\varphi}}, \tag{40}$$

we obtain the eigenmodes in the new coordinate system ($\tilde{t}$, $\tilde{\varphi}$) as follows

$$n_R = \frac{\Omega_H}{\bar{\varepsilon}+\varepsilon}[\omega - m(\Omega_H + \varepsilon)], \tag{41}$$

$$n_L = \frac{\omega - m\bar{\Omega}}{\alpha(\bar{\varepsilon}+\varepsilon)}. \tag{42}$$

In terms of these variables, the Boltzmann factor can be written as

$$e^{-(\omega - m\Omega_H)/T_H} = e^{-n_R/T_R - n_L/T_L}, \tag{43}$$

where the dimensionless left and right temperatures are given by

$$T_R = \frac{\Omega_H T_H}{\bar{\varepsilon}}, \tag{44}$$

$$T_L = \frac{T_H}{\alpha\varepsilon}, \tag{45}$$

with $T_H = \kappa_H/2\pi$.

Now, we calculate the entropy with the help of the Cardy formula. From Eqs. (33) and (44), the entropy of the right part becomes

$$S_R = \frac{\pi^2}{3} c_R T_R = \frac{A}{8G} \cdot \frac{4\Omega_H \bar{\varepsilon}^2}{q^{\varphi\varphi}\kappa_H^2 \rho^2}. \tag{46}$$

By setting $\bar{\varepsilon} = \frac{1}{2}\kappa_H \rho \sqrt{q^{\varphi\varphi}/\Omega_H}$, we obtain $S_R = A/(8G)$. The entropy of the left part can be obtained from Eqs. (39) and (45) as

$$S_L = \frac{\pi^2}{3} c_L T_L = \frac{A}{8G} \cdot \frac{4}{\alpha}, \tag{47}$$

which becomes $S_L = A/(8G)$ when $\alpha = 4$. Thus, by setting $\alpha = 4$ we obtain the total entropy of the system,

$$S = S_R + S_L = \frac{A}{4G}, \tag{48}$$

that agrees with the Bekenstein-Hawking entropy.

IV. CONCLUSION

We calculate separately the central charges c_L and c_R and the temperatures T_L and T_R on a stretched horizon of a nonextremal rotating black hole in the framework of Hamiltonian gravity. The entropy is obtained using the Cardy formula and it agrees with the Bekenstein-Hawking entropy $S_{\rm BH} = \frac{A}{4G}$, when the stretched horizon is set with $\bar{\varepsilon} = \frac{1}{2}\kappa_H \rho \sqrt{q^{\varphi\varphi}/\Omega_H}$ and the proportionality constant of the newly defined coordinate $\tilde{\varphi}$ is chosen appropriately.

In Refs. [45, 46], $\bar{\varepsilon}$ has been chosen such that the Killing vector $\bar{\chi}^\mu = \partial_t^\mu + \bar{\Omega}\partial_\varphi^\mu$ is null on the stretched horizon. This was done by imposing the null condition of the new Killing vector up to the order ρ^2:

$$\begin{aligned} 0 = \bar{\chi}^2 &= -N^2 + q_{\varphi\varphi}(N^\varphi + \bar{\Omega})^2 \\ &= -\kappa_H^2 \rho^2 + q_{\varphi\varphi}^{(H)} \bar{\varepsilon}^2 + 2 q_{\varphi\varphi}^{(H)} \bar{\varepsilon}\varepsilon + O(\rho^4). \end{aligned} \tag{49}$$

Namely, $\bar{\varepsilon}$ is set to satisfy $\bar{\varepsilon}^2 = \kappa_H^2 \rho^2 / q_{\varphi\varphi}^{(H)}$. However, if we do not restrict ourselves to the second order, then this condition does not satisfy the null condition, since $\bar{\varepsilon}$ and ε are the order of ρ and ρ^2, respectively.

In this paper, we do not impose the null condition for the Killing vector. Instead we try to get two Killing vectors each of which depends on one of the two independent coordinates separated slightly from the angular coordinate of the corotating frame at the horizon. This was done in the spirit of the point-splitting, but unlike the symmetric separations in the usual point-splitting method they are separated asymmetrically: one is the order of ρ and the other is much smaller, the order of ρ^2. We then obtain the central charges of the two Virasoro algebras resulting from these two Killing vectors. The two central charges and the Cardy formula yield correct Bekenstein-Hawking entropy once we set the parameters appropriately.

Finally, we make a note that when the two coordinates are separated equally, we could not get the correct Bekenstein-Hawking entropy. At this moment, we do not know the reason, but we speculate that asymmetric separations may play a role in keeping the two split coordinates independent. We leave this issue for future investigation.

Acknowledgments

ECY would like to thank KIAS for hospitality during the time that this work was done. This work was supported by National Research Foundation(NRF) of Korea Grants funded by the Korean Government (Ministry of Education, Science and Technology), NRF-2010-359-C00007(ME) and NRF-2011-0025517(ECY).

[1] M. Guica, T. Hartman, W. Song and A. Strominger, *The Kerr/CFT Correspondence, Phys. Rev.* D **80** (2009) 124008 [arXiv:0809.4266 [hep-th]].

[2] Y. Matsuo, T. Tsukioka and C. -M. Yoo, *Yet Another Realization of Kerr/CFT Correspondence, Europhys. Lett.* **89** (2010) 60001 [arXiv:0907.4272 [hep-th]].

[3] Y. Matsuo, T. Tsukioka and C. -M. Yoo, *Another Realization of Kerr/CFT Correspondence, Nucl. Phys.* B **825** (2010) 231 [arXiv:0907.0303 [hep-th]].

[4] I. Bredberg, C. Keeler, V. Lysov and A. Strominger, *Cargese Lectures on the Kerr/CFT Correspondence, Nucl. Phys. Proc. Suppl.* **216** (2011) 194 [arXiv:1103.2355 [hep-th]].

[5] B. Chen, B. Ning and J. -J. Zhang, *Boundary Conditions for NHEK through Effective Action*

Approach, Chin. Phys. Lett. **29** (2012) 041101 [arXiv:1105.2878 [hep-th]].

[6] G. Compere and S. Detournay, *Semi-classical central charge in topologically massive gravity, Class. Quant. Grav.* **26** (2009) 012001 [Erratum-ibid. **26** (2009) 139801] [arXiv:0808.1911 [hep-th]].

[7] G. Compere and S. Detournay, *Boundary conditions for spacelike and timelike warped AdS_3 spaces in topologically massive gravity, JHEP* **0908** (2009) 092 [arXiv:0906.1243 [hep-th]].

[8] M. Blagojevic and B. Cvetkovic, *Asymptotic structure of topologically massive gravity in spacelike stretched AdS sector, JHEP* **0909** (2009) 006 [arXiv:0907.0950 [gr-qc]].

[9] T. Hartman, K. Murata, T. Nishioka and A. Strominger, *CFT Duals for Extreme Black Holes, JHEP* **0904** (2009) 019 [arXiv:0811.4393 [hep-th]].

[10] G. Compere, K. Murata and T. Nishioka, *Central Charges in Extreme Black Hole/CFT Correspondence, JHEP* **0905** (2009) 077 [arXiv:0902.1001 [hep-th]].

[11] C. -M. Chen, J. -R. Sun and S. -J. Zou, *The RN/CFT Correspondence Revisited, JHEP* **1001** (2010) 057 [arXiv:0910.2076 [hep-th]].

[12] J. -J. Peng and S. -Q. Wu, *Extremal Kerr/CFT correspondence of five-dimensional rotating (charged) black holes with squashed horizons, Nucl. Phys.* B **828** (2010) 273 [arXiv:0911.5070 [hep-th]].

[13] M. R. Garousi and A. Ghodsi, *The RN/CFT Correspondence, Phys. Lett.* B **687** (2010) 79 [arXiv:0902.4387 [hep-th]].

[14] R. Li, M. -F. Li and J. -R. Ren, *Entropy of Kaluza-Klein Black Hole from Kerr/CFT Correspondence, Phys. Lett.* B **691** (2010) 249 [arXiv:1004.5335 [hep-th]].

[15] Y. Matsuo and T. Nishioka, *New Near Horizon Limit in Kerr/CFT, JHEP* **1012** (2010) 073 [arXiv:1010.4549 [hep-th]].

[16] M. Guica and A. Strominger, *Microscopic Realization of the Kerr/CFT Correspondence, JHEP* **1102** (2011) 010 [arXiv:1009.5039 [hep-th]].

[17] J. Rasmussen, *A near-NHEK/CFT correspondence, Int. J. Mod. Phys.* A **25** (2010) 5517 [arXiv:1004.4773 [hep-th]].

[18] M. Henneaux, C. Martinez and R. Troncoso, *Asymptotically warped anti-de Sitter spacetimes in topologically massive gravity, Phys. Rev.* D **84** (2011) 124016 [arXiv:1108.2841 [hep-th]].

[19] K. Goldstein and H. Soltanpanahi, *CFT Duals of Black Rings With Higher Derivative Terms, Class. Quant. Grav.* **29** (2012) 085016 [arXiv:1108.4362 [hep-th]].

[20] T. Azeyanagi, N. Ogawa and S. Terashima, *On Non-Chiral Extension of Kerr/CFT, JHEP* **1106** (2011) 081 [arXiv:1102.3423 [hep-th]].

[21] A. Guneratne, L. Rodriguez, S. Wickramasekara and T. Yildirim, *A non-Extremal Kerr-like AdS2xS2 Solution and Black-Hole/Near-Horizion-CFT Duality,* arXiv:1206.2261 [hep-th].

[22] B. K. Button, L. Rodriguez and C. A. Whiting, *A Near Horizon CFT Dual for Kerr-Newman-AdS, Int. J. Mod. Phys.* A **26** (2011) 3077 [arXiv:1009.1661 [hep-th]].

[23] L. Rodriguez and T. Yildirim, *Entropy and Temperature From Black-Hole/Near-Horizon-CFT Duality, Class. Quant. Grav.* **27** (2010) 155003 [arXiv:1003.0026 [hep-th]].

[24] I. Bredberg, T. Hartman, W. Song and A. Strominger, *Black Hole Superradiance From Kerr/CFT, JHEP* **1004** (2010) 019 [arXiv:0907.3477 [hep-th]].

[25] M. Cvetic and F. Larsen, *Greybody Factors and Charges in Kerr/CFT, JHEP* **0909** (2009) 088 [arXiv:0908.1136 [hep-th]].

[26] T. Hartman, W. Song and A. Strominger, *Holographic Derivation of Kerr-Newman Scattering Amplitudes for General Charge and Spin, JHEP* **1003** (2010) 118 [arXiv:0908.3909 [hep-th]].

[27] C. -M. Chen, Y. -M. Huang and S. -J. Zou, *Holographic Duals of Near-extremal Reissner-Nordstrom Black Holes, JHEP* **1003** (2010) 123 [arXiv:1001.2833 [hep-th]].

[28] A. Castro, A. Maloney and A. Strominger, *Hidden Conformal Symmetry of the Kerr Black Hole, Phys. Rev.* D **82** (2010) 024008 [arXiv:1004.0996 [hep-th]].

[29] R. Fareghbal, *Hidden Conformal Symmetry of the Warped* AdS_3 *Black Holes, Phys. Lett.* B **694** (2010) 138 [arXiv:1006.4034 [hep-th]].

[30] M. R. Setare and V. Kamali, *Twofold hidden conformal symmetries of rotating charged Goedel black holes, Class. Quant. Grav.* **28** (2011) 155006.

[31] C. -M. Chen and J. -R. Sun, *Holographic Dual of the Reissner-Nordstróm Black Hole, J. Phys. Conf. Ser.* **330** (2011) 012009 [arXiv:1106.4407 [hep-th]].

[32] B. Chen and J. -j. Zhang, *General Hidden Conformal Symmetry of 4D Kerr-Newman and 5D Kerr Black Holes, JHEP* **1108** (2011) 114 [arXiv:1107.0543 [hep-th]].

[33] E. Franzin and I. Smolic, *A New look at hidden conformal symmetries of black holes, JHEP* **1109** (2011) 081 [arXiv:1107.2756 [hep-th]].

[34] D. A. Lowe and A. Skanata, *Generalized Hidden Kerr/CFT,* arXiv:1112.1431 [hep-th].

[35] S. Bertini, S. L. Cacciatori and D. Klemm, *Conformal structure of the Schwarzschild black hole, Phys. Rev.* D **85** (2012) 064018 [arXiv:1106.0999 [hep-th]].

[36] M. R. Setare and V. Kamali, *Hidden conformal symmetry of rotating NS5-branes in extremal limit, Int. J. Mod. Phys.* A **26** (2011) 2233.

[37] M. R. Setare and V. Kamali, *Holographic description of a Kerr-Goedel black hole in five dimensions, Int. J. Mod. Phys.* A **26** (2011) 4287.

[38] C. -M. Chen, J. -R. Sun and J. -R. Sun, *The Kerr-Newman/CFTs Correspondence,* arXiv:1201.4040 [hep-th].

[39] A. M. Ghezelbash and H. M. Siahaan, *Hidden and Generalized Conformal Symmetry of Kerr-Sen Spacetimes,* arXiv:1206.0714 [hep-th].

[40] M. Cvetic and F. Larsen, *Conformal Symmetry for General Black Holes, JHEP* **1202** (2012) 122 [arXiv:1106.3341 [hep-th]].

[41] O. J. C. Dias, J. E. Santos and M. Stein, *Kerr-AdS and its Near-horizon Geometry: Perturbations and the Kerr/CFT Correspondence,* arXiv:1208.3322 [hep-th].

[42] S. Carlip, *Black hole entropy from conformal field theory in any dimension, Phys. Rev. Lett.* **82** (1999) 2828 [hep-th/9812013].

[43] M. -I. Park and J. Ho, *Comments on 'Black hole entropy from conformal field theory in any dimension', Phys. Rev. Lett.* **83** (1999) 5595 [hep-th/9910158].

[44] M. -I. Park, *Hamiltonian dynamics of bounded space-time and black hole entropy: Canonical method, Nucl. Phys.* B **634** (2002) 339 [hep-th/0111224].

[45] S. Carlip, *Extremal and nonextremal Kerr/CFT correspondences, JHEP* **1104** (2011) 076 [Erratum-ibid. **1201** (2012) 008] [arXiv:1101.5136 [gr-qc]].

[46] S. Carlip, *Effective Conformal Descriptions of Black Hole Entropy, Entropy* **13** (2011) 1355 [arXiv:1107.2678 [gr-qc]].

[47] B. Chen and J. -j. Zhang, *Novel CFT Duals for Extreme Black Holes, Nucl. Phys.* B **856** (2012) 449 [arXiv:1106.4148 [hep-th]].

[48] J. Mei, *On the General Kerr/CFT Correspondence in Arbitrary Dimensions, JHEP* **1204** (2012) 113 [arXiv:1202.4156 [hep-th]].

[49] J. D. Brown, S. R. Lau and J. W. York, Jr., *Action and energy of the gravitational field,* gr-qc/0010024.

[50] J. Novotny and M. Schnabl, *Point - splitting regularization of composite operators and anomalies, Fortsch. Phys.* **48** (2000) 253 [hep-th/9803244]; the references there in.